

A whale shark feeding in association with a school of giant herring at Ningaloo Reef, Western Australia

S G Wilson

Department of Zoology, The University of Western Australia, 35 Stirling Highway, Crawley WA 6009
Present address: Edgerton Research Laboratory, New England Aquarium, Central Wharf, Boston, MA 02110, USA
email: swilson@neaq.org

(Manuscript received November 2001; accepted February 2002)

Abstract. A whale shark was observed feeding among a school of giant herring at Ningaloo Reef. It is suggested that both the whale shark and the herring school were targeting the same planktonic prey.

Key words: whale shark, *Rhincodon typus*, giant herring, *Elops hawaiiensis*

Introduction

The whale shark, *Rhincodon typus*, is a suction filter-feeder that targets dense concentrations of planktonic and nektonic prey, including aggregating crustaceans such as euphausiids and copepods, and schooling baitfishes such as sardines and anchovies (Compagno 1984). Fishermen often use its presence as an indicator of waters rich in plankton and plankton-feeding fish that may, in turn, attract more valuable predatory species (Colman 1997). There are a number of reports of feeding associations occurring between whale sharks and schools of predatory fishes, such as mackerel, jacks and tuna (e.g. Gudger 1941; Springer 1957; Hoffman *et al.* 1981). In each of these cases, the whale sharks appeared to be targeting the same prey as the predatory fishes rather than the predatory fishes themselves.

Similar feeding associations have not been reported in whale sharks seasonally aggregating in coastal waters off Ningaloo Reef, Western Australia, where they are thought to feed predominantly on the neritic euphausiid *Pseudeuphausia latifrons*. On a number of occasions whale sharks have been observed feeding in these waters on surface schools of this species (Taylor & Grigg 1991; Clark 1992; Taylor 1994). Additionally, two whale shark faecal samples collected off Ningaloo Reef were found to contain crustacean remains resembling *P. latifrons* (Wilson & Newbound 2001). Other potential prey includes the large schools of baitfishes frequently sighted at this location at the time of the whale shark aggregation (Taylor 1994).

Materials and Methods

The author accompanied pilots of fixed-wing aircraft flown along the northern section of Ningaloo Reef from 4-11 May 2001.

Results

On 7 May 2001 at 1100 hrs, a whale shark was observed from the aircraft swimming northwards along the reef

front ~1 km south of South Passage (see Fig 1 in Wilson *et al.* 2001). A school of giant herring (*Elops hawaiiensis*), recognisable by their distinctive body shape, was visible feeding at the surface ~300 m to the northeast of the whale shark. Although not initially swimming on an intercept course, the shark made a 45-degree turn towards the school at a distance of ~100 m. Fig 1 shows the whale shark approaching the herring school. The shark stopped swimming when in the middle of the school (Fig 2), then adopted a stationary feeding posture (Fig 3) similar to that described by Springer (1957), Hoffman *et al.* (1981), Taylor *et al.* (1983) and Silas (1986), with its head just below the surface and its tail pointing downwards at a 45-degree angle. It remained in this



Figure 1. Whale shark approaches the herring school.



Figure 2. Whale shark stops swimming in the centre of the herring school.



Figure 3. Whale shark adopts a stationary feeding posture.

posture for a period of 3-4 minutes, after which time the herring school dispersed and the shark continued swimming northwards.

Discussion

It seems likely that the whale shark described here was targeting the same prey as the herring, rather than attempting to feed on the herring themselves. A whale shark tour operator reported a similar observation that took place in shallow water directly off the front of Ningaloo Reef (P O'Halloran, personal communication): the whale shark was hanging vertically in the water, with its head just below the surface and its tail sweeping along the sandy bottom, in the middle of a herring school; small, planktonic prey were being consumed by both the whale shark and the herring. This suggests that occurrences of whale sharks feeding among schools of giant herring may not be uncommon in these waters.

Acknowledgements: I would like to thank to P Arcott and the pilots of Exmouth Air Charters for their assistance in this work. This study was funded by Woodside Energy Ltd.

References

- Colman J (1997) Whale shark interaction management, with particular reference to Ningaloo Marine Park: 1997-2007. Western Australian Department of Conservation and Land Management, Western Australian Wildlife Management Program No. 27, 63 pp.
- Clark E 1992 Whale sharks: gentle monsters of the deep. *National Geographic* 182(6):123-138.
- Compagno L J V 1984 Sharks of the world, Part 1: Hexanchiformes to Lamniformes. FAO Fisheries Synopsis 125, Rome.
- Gudger E W 1941 The food and feeding habits of the whale shark, *Rhincodon typus*. *Journal of the Elisha Mitchell Scientific Society* 57:57-72.
- Hoffman W, Fritts T H & Reynolds R P 1981 Whale sharks associated with fish schools off south Texas. *Northeast Gulf Science* 5:55-56.
- Silas E G 1986 The whale shark (*Rhiniodon typus* Smith) in Indian coastal waters: is the species endangered or vulnerable? *Marine Fisheries Information Service, Technical and Extension Series* 66:1-19.
- Springer S 1957 Some observations on the behavior of schools of fishes in the Gulf of Mexico. *Ecology* 38:166-171.
- Taylor G 1994 Whale sharks: the giants of Ningaloo Reef. Angus & Robertson, Sydney.
- Taylor G & Grigg G 1991 Whale sharks of Ningaloo Reef. Australian National Parks and Wildlife Service, Commonwealth of Australia, Canberra, unpublished report, 20 pp.
- Taylor L R, Compagno L J V & Struhsaker P J 1983 Megamouth - a new species, genus, and family of lamnoid shark (*Megachasma pelagios*, Family Megachasmidae) from the Hawaiian Islands. *Proceedings of the California Academy of Sciences* 43:87-110.
- Wilson S G & Newbound D R 2001 Two whale shark faecal samples from Ningaloo Reef, Western Australia. *Bulletin of Marine Science* 68:361-362.
- Wilson S G, Pauly T & Meekan M G 2001 Daytime surface swarming by *Pseudeuphausia latifrons* (Crustacea, Euphausiacea) off Ningaloo Reef, Western Australia. *Bulletin of Marine Science* 68:157-162.